


ACTION: (check appropriate boxes)

- School Nurse to proceed directly to the symptomatic patient, with emergency medicine in hand.
- For accidental ingestion or exposure to _____ or stung by _____.
- Immediately give ____ mg. Benadryl (Diphenhydramine Hydrochloride).
- For any accidental exposure/sting - immediately administer epi-pen to outer portion of thigh, and dial 911. Inform 911 epinephrine may be needed. Transport to emergency room by ambulance.
- Student is self directed and may self carry emergency medication (as determined by MD and School Nurse).


DO NOT HESITATE TO ADMINISTER MEDICATION OR CALL RESCUE SQUAD

EPIPEN® AND EPIPEN® JR. DIRECTIONS

1. Pull off gray safety cap



2. Place black tip on outer thigh (always apply to thigh)



3. Using a swing and jab motion, press hard into thigh until Auto-Injector mechanism functions. Hold in place and count to 10. The EpiPen® unit should then be removed and taken with you to the Emergency Room. Massage the injection area for 10 seconds.

For children with multiple food allergies, use one form for each food.

MD Signature: _____ Date: _____ or see scripts

Printed MD Name _____ Or see attached scripts

Parent Signature: _____ Date: _____

School Nurse: _____ Date: _____

Must be updated/received yearly.